

SRI International

Final Report--Objective E, Task 1  
Covering the Period 1 October 1985 to 30 September 1987

December 1987

**POSSIBLE PHOTON PRODUCTION DURING  
A REMOTE VIEWING TASK:  
A REPLICATION EXPERIMENT (U)**

By:



**MANAGING TOTAL QUALITY**  
3M QUALITY MANAGEMENT SERVICES  
612/778-7560

*SRI Documents that have been  
declassified but PO has no classified  
originals so cannot create <sup>complete</sup> unclass.  
documents.*

Approved by:

*copy 2 of 10... copies.*

*This document consists of 19 pages.*

*SRI GE-0302*

*copy 1 of 3*



ABSTRACT (U)

Attempting to verify a claim by the Chinese that light is emitted in the vicinity of correctly identified remote viewing (RV) target material, we repeated an experiment first published in FY 1984. In that earlier experiment, a state-of-the-art, ambient temperature, photon counting system was used to monitor the target material (35-mm slides of *National Geographic Magazine* photographs). The statistical measure derived from the photon counting apparatus in that study showed a significant positive correlation with the RV results ( $p \leq 0.035$ ). That is, when the remote viewing was good, there was an increase in the signal detected by the photon counting system. In addition, we observed two anomalous pulses having a signal-to-noise ratio of about 20:1 to 40:1. In the present experiment (FY 1987), we improved all hardware aspects of the previous work, substantially reducing the background noise level and improving shielding against artifact. In addition, analysis of the remote viewing indicates that three out of the four viewers produced independently significant results. Our analysis of the photomultiplier tube (PMT) data shows no evidence of any anomalous high count rate pulses, no evidence of any effect on the PMT output during the RV session, and no evidence of any significant correlation between RV performance and PMT output. We conclude that (1) the effect proposed by the Chinese is artifactual in nature.

## III RESULTS (U)

## A. (U) Remote Viewing Results

(U) Each RV session was judged using an FM analysis. The FM is defined as the product of two measures: accuracy and reliability. The accuracy of an RV response is the fraction of the target material that is described correctly. Reliability is the fraction of the response that is correct. Tables I through 4 show the RV results for each trial. The session number (9001.cr, for example) incorporates a code for each viewer as well as the chronological sequence of viewings.

Table 1

(U) REMOTE VIEWING RESULTS FOR VIEWER 009

Session	Figure of Merit	p-value
9001.lg	0.5714	0.0238
9002.lg	0.3810	0.1961
9003.lg	0.4444	0.0497
9004.lg	0.3333	0.3650
9005.lg	0.0667	0.9233
9006.lg	0.3556	0.2697
Overall $p \leq 0.0450$		

Table 2

(U) REMOTE VIEWING RESULTS FOR VIEWER 105

Session	Figure of Merit	p-value
9001.rs	0.4571	0.0412
9002.rs	0.1667	0.3486
9003.rs	0.1600	0.3618
9004.rs	0.3333	0.1039
9005.rs	0.0000	1.0000
9006.rs	0.3810	0.0475
Overall $p \leq 0.0488$		

Table 3

(U) REMOTE VIEWING RESULTS FOR VIEWER 177

Session	Figure of Merit	p-value
9001.hs	0.4444	0.2430
9002.hs	0.1143	0.9579
9003.hs	0.3810	0.2978
9004.hs	0.5000	0.2392
9005.hs	0.5952	0.0677
9006.hs	0.6429	0.0136
Overall $p \leq 0.0385$		

Table 4

(U) REMOTE VIEWING RESULTS FOR VIEWER 807

Session	Figure of Merit	p-value
9001.cr	0.0000	1.0000
9002.cr	0.3333	0.2267
9003.cr	0.5208	0.0240
9004.cr	0.0833	0.7494
9005.cr	0.3750	0.1321
9006.cr	0.1333	0.5911
Overall $p \leq 0.1895$ , n.s.		

(U) From the FM analysis performed for our FY 1984 experiment, we determined that by computing the p-value for each FM we could determine an average p for each viewer and for all sessions combined. The overall probability of obtaining that average p-value was then calculated, either by an exact method for small numbers of sessions<sup>7</sup> or by using the central limit theorem for greater than 20 sessions.<sup>8</sup> In the current analysis, an additional test of significance, the Fisher Chi-square technique,<sup>8</sup> has been added to supplement the probability associated with average p-value for a given series.

The overall p-values given for each viewer's series as shown in Tables 1 through 4 were calculated using the Fisher Chi-square technique. Averaging all p-values for all sessions yielded  $p(\text{avg.}) = 0.3437$ . Using the central-limit theorem, the probability associated with that average value is  $p \leq 0.004$ . Using the Fisher Chi-square method, a p-value of  $\leq 0.0036$  was calculated for all 24 sessions, indicating good agreement between techniques. We observed that three out of the four viewers independently produced significant results. Such an outcome is an extremely rare event. If the probability of success is  $p \leq 0.05$ , the binomial probability of obtaining three out of four successful results is  $p \leq 0.00048$ . These individual and overall remote viewing results are substantially better than were achieved in the FY 1984 study.